

Dissipative properties of the cosmological fluid

Stefan Flörchinger

Max-Planck Institut für Kernphysik, Heidelberg, 11. Juli 2016.





UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386

Content

- Why are dissipative properties of dark matter interesting ?
- How to understand large scale structure formation ?
- Could dissipation affect the cosmological expansion ?
- How do dissipation and thermalization work in general relativity ?

Why are dissipative properties of dark matter interesting ?

Big bang – little bang analogy



Cosmology

- cosmol. scale: $\text{Mpc} = 3.1 \times 10^{22} \text{ m}$
- Gravity + QED + Dark sector
- one big event

Heavy ion collisions

- nuclear scale: $\text{fm} = 10^{-15} \text{ m}$
- QCD
- very many events
- initial conditions not directly accessible
- all information must be reconstructed from final state
- **dynamical description as a (viscous) fluid**

The dark matter fluid

- Heavy ion collisions

$$\mathcal{L}_{\text{QCD}} \rightarrow \text{fluid properties}$$

- Late time cosmology

$$\text{fluid properties} \rightarrow \mathcal{L}_{\text{dark matter}}$$

- Until direct detection of dark matter, it can only be observed via

$$T_{\text{dark matter}}^{\mu\nu}$$

Energy-momentum tensor and conserved current


$$T^{\mu\nu} = (\epsilon + p + \pi_{\text{bulk}})u^\mu u^\nu + (p + \pi_{\text{bulk}})g^{\mu\nu} + \pi^{\mu\nu}$$

$$N^\mu = n u^\mu + \nu^\mu$$

- tensor decomposition w. r. t. fluid velocity u^μ
- pressure $p = p(\epsilon, n)$
- constitutive relations for viscous terms in derivative expansion
 - bulk viscous pressure $\pi_{\text{bulk}} = -\zeta \nabla_\mu u^\mu + \dots$
 - shear stress $\pi^{\mu\nu} = -\eta \left[\Delta^{\mu\alpha} \nabla_\alpha u^\nu + \Delta^{\nu\alpha} \nabla_\alpha u^\mu - \frac{2}{3} \Delta^{\mu\nu} \nabla_\alpha u^\alpha \right] + \dots$
 - diffusion current $\nu^\alpha = -\kappa \left[\frac{nT}{\epsilon + p} \right]^2 \Delta^{\alpha\beta} \partial_\beta \left(\frac{\mu}{T} \right) + \dots$

Fluid dynamic equations from covariant **conservation laws**

$$\nabla_\mu T^{\mu\nu} = 0, \quad \nabla_\mu N^\mu = 0.$$



- Long distances, long times or strong enough interactions
- Needs **macroscopic** fluid properties
 - equation of state $p(\epsilon, n)$
 - shear viscosity $\eta(\epsilon, n)$
 - bulk viscosity $\zeta(\epsilon, n)$
 - heat conductivity $\kappa(\epsilon, n)$
 - relaxation times, ...
- For QCD no full *ab initio* calculation of transport properties possible yet but in principle fixed by **microscopic** properties encoded in \mathcal{L}_{QCD}
- Ongoing experimental and theoretical effort to understand this in detail

Ideal fluid versus collision-less gas

- Many codes used in cosmology describe dark matter as **ideal, cold and pressure-less fluid**

$$T^{\mu\nu} = \epsilon u^{\mu} u^{\nu}$$

- Equation of state $p = 0$
- No shear stress and bulk viscous pressure $\pi^{\mu\nu} = \pi_{\text{bulk}} = 0$
- Dark matter is also modeled as **collision-less gas** of massive particles, interacting via gravity only
- Two pictures are in general **not consistent**

Dissipative properties


Viscosities

- Diffusive transport of momentum [Maxwell (1860)]
- Depend strongly on interaction properties
- Example: non-relativistic gas of particles with mass m , mean peculiar velocity \bar{v} , elastic $2 \rightarrow 2$ cross-section σ_{el}

$$\eta = \frac{m \bar{v}}{3 \sigma_{\text{el}}} \quad \zeta = 0$$

- Interesting additional **information about dark matter**

Self-interaction of dark matter





Gravitational lensing and x-ray image of “bullet cluster” 1E0657-56

- so far: dark matter is non-interacting → can collide without stopping

$$\frac{\sigma_{\text{el}}}{m} \lesssim 1.2 \frac{\text{cm}^2}{\text{g}}$$

Is dark matter self-interacting?



Galaxy cluster Abell 3827

[Massey *et al.*, MNRAS 449, 3393 (2015)]

- Offset between stars and dark matter falling into cluster
- Is this a first indication for a dark matter self-interaction?

$$\frac{\sigma_{\text{el}}}{m} \approx 3 \frac{\text{cm}^2}{\text{g}} \approx 0.5 \frac{\text{b}}{\text{GeV}} \quad (\text{under debate})$$

[Kahlhoefer, Schmidt-Hoberg, Kummer & Sarkar, MNRAS 452, 1 (2015)]

Precision cosmology can measure shear stress

- Scalar excitations in gravity

$$ds^2 = a^2 [-(1 + 2\psi)d\eta^2 + (1 - 2\phi)dx_i dx_i]$$

with two Newtonian potentials ψ and ϕ .

- Einsteins equations imply

$$(\partial_i \partial_j - \frac{1}{3} \delta_{ij} \partial_k^2) (\phi - \psi) = 8\pi G_N a^2 \pi_{ij}|_{\text{scalar}}$$


with scalar part of shear stress

$$\pi_{ij}|_{\text{scalar}} = (\partial_i \partial_j - \frac{1}{3} \delta_{ij} \partial_k^2) \tilde{\pi}$$

- Detailed data at small redshift e.g. from Euclid satellite (esa, 2020)
[Amendola *et al.* (2012)]
 - ψ can be measured via acceleration of matter
 - $\psi + \phi$ can be measured by weak lensing and Sachs-Wolfe effect
 - fluid velocity can be accessed by redshift space distortions
- New quantitative precise insights into fluid properties of dark matter

Cosmological structure formation

- How do **viscosities influence structure formation**?
- Does viscous fluid dynamics help to **understand large scale structure (semi) analytically**?



[Springel, Frenk & White,
Nature 440, 1137 (2006)]

How is structure formation modified?

Linear dynamics

- energy conservation ($\theta = \vec{\nabla} \cdot \vec{v}$)

$$\dot{\delta\epsilon} + 3\frac{\dot{a}}{a}\delta\epsilon + \bar{\epsilon}\theta = 0$$

- Navier-Stokes equation

$$\bar{\epsilon} \left[\dot{\theta} + \frac{\dot{a}}{a}\theta - k^2\psi \right] + \frac{1}{a} \left(\zeta + \frac{4}{3}\eta \right) k^2\theta = 0$$

- Poisson equation


$$-k^2\psi = 4\pi G_N a^2 \delta\epsilon$$

Scalar perturbations ($\delta = \frac{\delta\epsilon}{\bar{\epsilon}}$)

$$\ddot{\delta} + \left[\frac{\dot{a}}{a} + \frac{\zeta + \frac{4}{3}\eta}{a\bar{\epsilon}} k^2 \right] \dot{\delta} - 4\pi G_N \bar{\epsilon} \delta = 0$$

Viscosities slow down gravitational collapse but do not wash out structure

Structure formation with viscosities



[Blas, Floerchinger, Garny, Tetradis & Wiedemann, JCAP 1511, 049 (2015)]

- k -dependent growth factor for scalar modes
- Could be tested by observation of large scale structure
- Depends on $\zeta + \frac{4}{3}\eta$ as function of time (or density)

How to understand large scale structure formation ?

Formation of large scale structure

- Formation of large scale structure is interesting
 - tests physics of dark matter
 - tests physics of dark energy
 - gets tested by missions like Euclid, ...
- Cosmological perturbation theory breaks down when density contrast

$$\delta(\mathbf{k}) = \frac{\delta\rho(\mathbf{k})}{\bar{\rho}} \gg 1$$

grows large at late times and for small scales.

- Numerical simulations (N -body) are expensive and time-consuming
- One would like to have better analytical understanding

Renormalization group approach

[Blas, Floerchinger, Garny, Tetradis & Wiedemann, JCAP 1511, 049 (2015)]

[Floerchinger, Garny, Tetradis & Wiedemann, in preparation]

- Start from ideal fluid approximation
- Large scale structure formation can be formulated as classical field theory with stochastic initial conditions
- Leads to classical statistical field theory
- Initial state fluctuations can be treated by functional renormalization group, similar to thermal or quantum fluctuations in other contexts [Matarrese & Pietroni (2007)]
- Modify theory by cutting off the initial spectrum in the IR

$$P_k^0(\mathbf{q}) = P^0(\mathbf{q}) \Theta(|\mathbf{q}| - k)$$

- Use flow equation for 1PI effective action [Wetterich (1993)]

$$\partial_k \Gamma_k[\phi, \chi] = \frac{1}{2} \text{Tr} \left\{ \left(\Gamma_k^{(2)}[\phi, \chi] - i (P_k^0 - P^0) \right)^{-1} \partial_k P_k^0 \right\}$$

Renormalization of effective viscosity and pressure

- Effective theory at scale k has additional terms in equations of motion
- Order them by derivative expansion.
- Lowest order: ideal fluid
- Next-to-lowest order: effective sound velocity parameter

$$\gamma_s = \frac{c_s^2}{\mathcal{H}^2} = \frac{dp/d\rho}{\mathcal{H}^2}.$$

and effective viscosity parameter



$$\gamma_\nu = \frac{4\eta/3 + \zeta}{(\rho + p)\mathcal{H}a}.$$

- Both depend on cosmological time or scale factor a

$$\gamma_s = \lambda_s a^\kappa, \quad \gamma_\nu = \lambda_\nu a^\kappa$$

with exponent $\kappa \approx 2$.

RG flow of effective sound velocity parameter





- left: RG flow of effective sound velocity today $\lambda_s = \frac{c_s^2}{H_0^2} = \frac{dp/d\rho}{H_0^2}$
 - dashed line: one-loop approximation

$$\partial_k \lambda_s = -\frac{4\pi}{3} \frac{31}{70} P^0(k)$$

- solid line: functional RG
- right: linear density power spectrum

RG flow of effective viscosity parameter





- left: RG flow of effective viscosity today $\lambda_\nu = \frac{4\eta/3+\zeta}{(\rho+p)_0 H_0}$
 - dashed line: one-loop approximation

$$\partial_k \lambda_\nu = -\frac{4\pi}{3} \frac{78}{35} P^0(k)$$

- solid line: functional RG
- right: linear density power spectrum

RG flow of exponent κ




- left: RG flow of exponent κ
 - dashed line: one-loop approximation

$$\partial_k \kappa = \frac{4\pi}{3} P^0(k) \frac{78(\kappa - 2)}{35\tilde{\lambda}_\nu}$$

- solid line: functional RG
- right: linear density power spectrum

Fixed point behavior



- growing mode is sensitive to $\lambda_s + \lambda_\nu$
- functional RG has IR fixed points


Functional RG + perturbation theory

[Blas, Floerchinger, Garny, Tetradis & Wiedemann, JCAP 1511, 049 (2015)]


[Floerchinger, Garny, Tetradis & Wiedemann, in preparation]

- RG evolution to determine effective viscosity and sound velocity at intermediate scale k_m
- Perturbation theory for power spectrum for scales $0 < |\mathbf{q}| < k_m$
- Theory with effective parameters

$P_{\delta\delta}(k, z=0)$, viscous theory



$P_{\delta\delta}(k, z=0)$, SPT with cutoff



Could dissipation affect the cosmological expansion ?

Bulk viscosity

- Bulk viscous pressure is negative for expanding universe

$$\pi_{\text{bulk}} = -\zeta \nabla_{\mu} u^{\mu} = -\zeta 3H < 0$$

- Negative effective pressure

$$p_{\text{eff}} = p + \pi_{\text{bulk}} < 0$$

would act similar to dark energy in Friedmann's equations

[Murphy (1973), Padmanabhan & Chitre (1987), Fabris, Goncalves & de Sa Ribeiro (2006), Li & Barrow (2009), Velten & Schwarz (2011), Gagnon & Lesgourgues (2011), ...]





- **Is negative effective pressure physical?**
- In context of heavy ion physics: instability for $p_{\text{eff}} < 0$ (“cavitation”)
[Torrieri & Mishustin (2008), Rajagopal & Tripuraneni (2010), Buchel, Camanho & Edelstein (2014), Habich & Romatschke (2015), Denicol, Gale & Jeon (2015)]
- What precisely happens at the instability?

Is negative effective pressure physical?

- Kinetic theory

$$p_{\text{eff}}(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{\vec{p}^2}{3E_{\vec{p}}} f(x, \vec{p}) \geq 0$$


- Stability argument



If there is a vacuum with $\epsilon = p_{\text{eff}} = 0$, phases with $p_{\text{eff}} < 0$ cannot be mechanically stable. (But could be metastable.)

Bulk viscosity in heavy ion physics

- In heavy ion physics people start now to consider bulk viscosity.
- Becomes relevant close to chiral crossover



[Denicol, Gale & Jeon (2015)]

- Is there a first-order phase transition triggered by the expansion?
- What is the relation to chemical and kinetic freeze-out?
- More detailed understanding needed, both for heavy ion physics and cosmology

Backreaction: General idea

- for 0 + 1 dimensional, non-linear dynamics

$$\dot{\varphi} = f(\varphi) = f_0 + f_1 \varphi + \frac{1}{2} f_2 \varphi^2 + \dots$$

- one has for expectation values $\bar{\varphi} = \langle \varphi \rangle$

$$\dot{\bar{\varphi}} = f_0 + f_1 \bar{\varphi} + \frac{1}{2} f_2 \bar{\varphi}^2 + \frac{1}{2} f_2 \langle (\varphi - \bar{\varphi})^2 \rangle + \dots$$

- evolution equation for expectation value $\bar{\varphi}$ depends on two-point correlation function or spectrum $P_2 = \langle (\varphi - \bar{\varphi})^2 \rangle$
- evolution equation for spectrum depends on bispectrum and so on
- more complicated for higher dimensional theories
- more complicated for gauge theories such as gravity

Backreaction in gravity

- Einstein's equations are non-linear.
- Important question [G. F. R. Ellis (1984)]: If Einstein's field equations describe small scales, including inhomogeneities, do they also hold on large scales?
- Is there a sizable backreaction from inhomogeneities to the cosmological expansion?
- Difficult question, has been studied by many people
[Ellis & Stoeger (1987); Mukhanov, Abramo & Brandenberger (1997); Unruh (1998); Buchert (2000); Geshnzjani & Brandenberger (2002); Schwarz (2002); Wetterich (2003); Räsänen (2004); Kolb, Matarrese & Riotto (2006); Brown, Behrend, Malik (2009); Gasperini, Marozzi & Veneziano (2009); Clarkson & Umeh (2011); Green & Wald (2011); ...]
- Recent reviews: [Buchert & Räsänen, Ann. Rev. Nucl. Part. Sci. 62, 57 (2012); Green & Wald, Class. Quant. Grav. 31, 234003 (2014)]
- No general consensus but most people believe now that **gravitational backreaction is rather small**.
- In the following we look at a new **backreaction on the matter side** of Einstein's equations.

Fluid equation for energy density

First order viscous fluid dynamics

$$u^\mu \partial_\mu \epsilon + (\epsilon + p) \nabla_\mu u^\mu - \zeta \Theta^2 - 2\eta \sigma^{\mu\nu} \sigma_{\mu\nu} = 0$$

For $\vec{v}^2 \ll c^2$ and Newtonian potentials $\Phi, \Psi \ll 1$

$$\begin{aligned} & \dot{\epsilon} + \vec{v} \cdot \vec{\nabla} \epsilon + (\epsilon + p) \left(3 \frac{\dot{a}}{a} + \vec{\nabla} \cdot \vec{v} \right) \\ &= \frac{\zeta}{a} \left[3 \frac{\dot{a}}{a} + \vec{\nabla} \cdot \vec{v} \right]^2 + \frac{\eta}{a} \left[\partial_i v_j \partial_i v_j + \partial_i v_j \partial_j v_i - \frac{2}{3} (\vec{\nabla} \cdot \vec{v})^2 \right] \end{aligned}$$

Fluid dynamic backreaction in Cosmology

[Floerchinger, Tetradis & Wiedemann, PRL 114, 091301 (2015)]

Expectation value of energy density $\bar{\epsilon} = \langle \epsilon \rangle$

$$\frac{1}{a} \dot{\bar{\epsilon}} + 3H (\bar{\epsilon} + \bar{p} - 3\bar{\zeta}H) = D$$

with dissipative backreaction term

$$D = \frac{1}{a^2} \langle \eta [\partial_i v_j \partial_i v_j + \partial_i v_j \partial_j v_i - \frac{2}{3} \partial_i v_i \partial_j v_j] \rangle \\ + \frac{1}{a^2} \langle \zeta [\vec{\nabla} \cdot \vec{v}]^2 \rangle + \frac{1}{a} \langle \vec{v} \cdot \vec{\nabla} (p - 6\zeta H) \rangle$$

- D vanishes for unperturbed homogeneous and isotropic universe
- D has contribution from shear & bulk viscous dissipation and thermodynamic work done by contraction against pressure gradients
- dissipative terms in D are positive semi-definite
- for spatially constant viscosities and scalar perturbations only

$$D = \frac{\bar{\zeta} + \frac{4}{3}\bar{\eta}}{a^2} \int d^3q P_{\theta\theta}(q)$$

Dissipation of perturbations

[Floerchinger, Tetradis & Wiedemann, PRL 114, 091301 (2015)]

- Dissipative backreaction does not need negative effective pressure

$$\frac{1}{a} \dot{\bar{\epsilon}} + 3H (\bar{\epsilon} + \bar{p}_{\text{eff}}) = D$$

- D is an integral over perturbations, could become large at late times.
- Can it potentially accelerate the universe?
- Need additional equation for scale parameter a
- Use trace of Einstein's equations $R = 8\pi G_{\text{N}} T^\mu{}_\mu$

$$\frac{1}{a} \dot{H} + 2H^2 = \frac{4\pi G_{\text{N}}}{3} (\bar{\epsilon} - 3\bar{p}_{\text{eff}})$$

does not depend on unknown quantities like $\langle (\epsilon + p_{\text{eff}}) u^\mu u^\nu \rangle$

- To close the equations one needs equation of state $\bar{p}_{\text{eff}} = \bar{p}_{\text{eff}}(\bar{\epsilon})$ and dissipation parameter D


Deceleration parameter

[Floerchinger, Tetradis & Wiedemann, PRL 114, 091301 (2015)]

- assume now vanishing effective pressure $\bar{p}_{\text{eff}} = 0$
- obtain for deceleration parameter $q = -1 - \frac{\dot{H}}{aH^2}$

$$-\frac{dq}{d \ln a} + 2(q - 1) \left(q - \frac{1}{2}\right) = \frac{4\pi G_{\text{N}} D}{3H^3}$$

- for $D = 0$ attractive fixed point at $q_* = \frac{1}{2}$ (deceleration)
- for $D > 0$ fixed point shifted towards $q_* < 0$ (acceleration)



Estimating viscous backreaction D

- For $\frac{4\pi G_N D}{3H^3} \approx 4$ one could explain the current accelerated expansion ($q \approx -0.6$) by dissipative backreaction.
- Is this possible?
- In principle one can determine D for given equation of state and viscous properties from dynamics of structure formation.
- So far only rough estimates. If shear viscosity dominates:

$$D = \frac{1}{a^2} \langle \eta [\partial_i v_j \partial_i v_j + \partial_i v_j \partial_j v_i - \frac{2}{3} \partial_i v_i \partial_j v_j] \rangle \approx c_D \bar{\eta} H^2$$

with $c_D = \mathcal{O}(1)$. Corresponds to $\Delta v \approx 100$ km/s for $\Delta x \approx 1$ MPc

- Leads to

$$\frac{4\pi G_N D}{3H^3} \approx \frac{c_D \bar{\eta} H}{2\rho_c}$$

with $\rho_c = \frac{3H^2}{8\pi G_N}$

Viscosities

- Relativistic particles / radiation contribute to shear viscosity

$$\eta = c_\eta \epsilon_R \tau_R$$

- prefactor $c_\eta = \mathcal{O}(1)$
- energy density of radiation ϵ_R
- mean free time τ_R
- Bulk viscosity vanishes in situations with conformal symmetry but can be large when conformal symmetry is broken.
- For massive scalar particles with $\lambda\varphi^4$ interaction [Jeon & Yaffe (1996)]

$$\zeta \sim \frac{m^6}{\lambda^4 T^3} e^{2m/T}, \quad \eta \sim \frac{m^{5/2} T^{1/2}}{\lambda^2} \quad \text{for} \quad \frac{T}{m} \ll 1$$

Estimating viscous backreaction D

Consider shear viscosity from radiation

$$\eta = c_\eta \epsilon_R \tau_R$$

Backreaction term

$$\frac{4\pi G_N D}{3H^3} \approx \frac{c_D c_\eta}{2} \frac{\epsilon_R}{\rho_c} \tau_R H$$

- fluid approximation needs $\tau_R H < 1$
- for sizeable effect one would need $\epsilon_R/\rho_c = \mathcal{O}(1)$
- unlikely that D becomes large enough in this scenario

Needed refinements:

- full dynamics of perturbations
- second order fluid dynamics
- complete model(s)

How do dissipation and thermalization work in general relativity ?

Gravity and thermalization

Consider ensemble of massive particles interacting via gravity only. Start with some velocity distribution. Is there **equilibration/thermalization**...

- ... in Newtonian gravity?
- ... in classical General relativity?
- ... in quantized gravity?

Analogy to other gauge theories suggests that **quantum properties are important for thermalization**

Dissipation by gravity

- Gravitational waves in viscous fluid have life time [Hawking (1966)]

$$\tau_G = \frac{1}{16\pi G_N \eta}$$

- Diffusive momentum transport by graviton radiation induces viscosity

$$\eta \approx \epsilon_G \tau_G$$

with energy density of gravitational field ϵ_G

- Can be solved for η and τ_G [Weinberg (1972)]

$$\eta = \sqrt{\frac{\epsilon_G}{16\pi G_N}}, \quad \tau_G = \sqrt{\frac{1}{16\pi G_N \epsilon_G}}$$

- Can this really be independent of dark matter mass and density?
- Thermalization time $\sim m_P/T^2$ is very large
- What determines dissipation on shorter time scales, when classical fields dominate?

Conclusions

Dissipative properties of the cosmological fluid...

- ... are quite interesting
- ... could provide more detailed understanding of dark matter
- ... can be tested by precision cosmology
- ... can help to better understand large scale structure
- ... could even affect the cosmological expansion

Backup slides

“Fundamental” and “effective” viscosity


Two types of viscosities for cosmological fluid

- ① Momentum transport by particles or radiation
 - governed by interactions
 - from linear response theory [Green (1954), Kubo (1957)]
 - close to equilibrium


- ② Momentum transport in the inhomogeneous, coarse-grained fluid
 - governed by non-linear fluid mode couplings
 - determined perturbatively [Blas, Floerchinger, Garny, Tetradis & Wiedemann]
 - non-equilibrium
 - heavy ions: anomalous plasma viscosity [Asakawa, Bass & Müller (2006)]
eddy viscosity [Romatschke (2008)]

Power spectrum at different redshifts


$P_{\delta\delta}(k, z=0), k_m = 0.6h/\text{Mpc}$




$P_{\delta\delta}(k, z=0.375), k_m = 0.6h/\text{Mpc}$



$P_{\delta\delta}(k, z=0.833), k_m = 0.6h/\text{Mpc}$




$P_{\delta\delta}(k, z=1.75), k_m = 0.6h/\text{Mpc}$




Velocity spectra


$P_{\theta\theta}(k, z=0)/P_{\delta\delta}$, viscous theory




$P_{\theta\theta}(k, z=0)/P_{\delta\delta}$, SPT with cutoff




$P_{\delta\theta}(k, z=0)/P_{\delta\delta}$, viscous theory



$P_{\delta\theta}(k, z=0)/P_{\delta\delta}$, SPT with cutoff



Power spectrum, standard perturbation theory



[D. Blas, M. Garny and T. Konstandin, JCAP 1309 (2013) 024]

Could viscous backreaction lead to Λ CDM-type expansion?

[Floerchinger, Tetradis & Wiedemann, 1506.00407]

- Backreaction term $D(z)$ will be *some* function of redshift.
- For given dissipative properties $D(z)$ can be determined, but calculation is involved.
- One may ask simpler question: For what form of $D(z)$ would the expansion be as in the Λ CDM model?
- The *ad hoc* ansatz $D(z) = \text{const} \cdot H(z)$ leads to modified Friedmann equations

$$\bar{\epsilon} - \frac{D}{4H} = \frac{3}{8\pi G_N} H^2, \quad \bar{p}_{\text{eff}} - \frac{D}{12H} = -\frac{1}{8\pi G_N} \left(2\frac{1}{a} \dot{H} + 3H^2 \right)$$

- In terms of $\hat{\epsilon} = \bar{\epsilon} - \frac{D}{3H}$ one can write

$$\frac{1}{a} \dot{\hat{\epsilon}} + 3H(\hat{\epsilon} + \bar{p}_{\text{eff}}) = 0, \quad R + \frac{8\pi G_N D}{3H} = -8\pi G_N (\hat{\epsilon} - 3\bar{p}_{\text{eff}})$$

- For $\bar{p}_{\text{eff}} = 0$ these are standard equations for Λ CDM model with

$$\Lambda = \frac{2\pi G_N D}{3H}$$

Modification of Friedmann's equations by backreaction 1

- For universe with fluid velocity inhomogeneities one cannot easily take direct average of Einstein's equations.
- However, fluid equation for energy density and trace of Einstein's equations can be used.
- By integration one finds modified Friedmann equation

$$H(\tau)^2 = \frac{8\pi G_N}{3} \left[\bar{\epsilon}(\tau) - \int_{\tau_1}^{\tau} d\tau' \left(\frac{a(\tau')}{a(\tau)} \right)^4 a(\tau') D(\tau') \right]$$

- Additive deviation from Friedmann's law for $D(\tau') > 0$
- Part of the total energy density is due to dissipative production

$$\bar{\epsilon} = \bar{\epsilon}_{\text{nd}} + \bar{\epsilon}_{\text{d}}$$

- Assume for dissipatively produced part

$$\dot{\bar{\epsilon}}_{\text{d}} + 3 \frac{\dot{a}}{a} (1 + \hat{w}_{\text{d}}) \bar{\epsilon}_{\text{d}} = a D$$

Modification of Friedmann's equations by backreaction 2

Leads to another variant of Friedmann's equation

$$H(\tau)^2 = \frac{8\pi G_N}{3} \left[\bar{\epsilon}_{\text{nd}}(\tau) + \int_{\tau_1}^{\tau} d\tau' \left[\left(\frac{a(\tau')}{a(\tau)} \right)^{3+3\hat{w}_d} - \left(\frac{a(\tau')}{a(\tau)} \right)^4 \right] a(\tau') D(\tau') \right]$$

- If the dissipative backreaction D produces pure radiation, $\hat{w}_d = 1/3$, it does not show up in effective Friedmann equation at all!
- For $\hat{w}_d < 1/3$ there is a new component with positive contribution on the right hand side of the effective Friedmann equation.
- To understand expansion, parametrize for late times

$$D(\tau) = H(\tau) \left(\frac{a(\tau)}{a(\tau_0)} \right)^{-\kappa} \tilde{D}$$

with constants \tilde{D} and κ .

- Hubble parameter as function of $(a_0/a) = 1 + z$

$$H(a) = H_0 \sqrt{\Omega_\Lambda + \Omega_M \left(\frac{a_0}{a} \right)^3 + \Omega_R \left(\frac{a_0}{a} \right)^4 + \Omega_D \left(\frac{a_0}{a} \right)^\kappa}$$

- For $\kappa \approx 0$ the role of Ω_Λ and Ω_D would be similar.

Inhomogeneities in heavy ion collisions


Inhomogeneities are main source of information in cosmology.

Similarly, in heavy ion collisions:

- **Initial fluid perturbations:** Event-by-event fluctuations around averaged fluid fields at time τ_0 and their evolution:
 - energy density ϵ
 - fluid velocity u^μ
 - shear stress $\pi^{\mu\nu}$
 - more general also: baryon number density n , electric charge density, electromagnetic fields, ...
- governed by universal evolution equations
- determine particle distributions after freeze-out, e.g. $v_n(p_T)$
- usefull to constrain **thermodynamic and transport properties of QCD**
- contain interesting information from early times

Fluid dynamic perturbation theory for heavy ions

[Floerchinger & Wiedemann, PLB 728, 407 (2014); JHEP 08 (2014) 005]



- goal: understand dynamics of heavy ion collisions better, constrain fluid properties of QCD from experimental results
- before: fully numerical fluid simulations e.g. [Heinz & Snellings (2013)]
- new: solve fluid equations for smooth and symmetric background and order-by-order in perturbations
- similar to cosmological perturbation theory
- good convergence properties [Floerchinger *et al.*, PLB 735, 305 (2014), Brouzakis *et al.* PRD 91, 065007 (2015)]
- some insights from heavy ion physics might be useful for cosmology

Fluid dynamics in cosmology and heavy ion collisions

Cosmology

- Large part of cosmological evolution is governed by equilibrium thermodynamics or ideal fluid dynamics.
- Free-streaming of photons and neutrinos at late times.
- Matter in equilibrium at early times, drops out of equilibrium later.
- Gravitational interaction is long range and treated explicitly.

Heavy ion collisions

- Expansion governed by viscous fluid dynamics.
- Free streaming of hadrons at late times.
- Strong interactions are confined at low T and screened at high T , treated implicitly.

First steps towards fluid dynamic perturbation or response theory

- Linear perturbations around Bjorken flow [Floerchinger & Wiedemann (2011)]
- Linear perturbations around Gubser solution for conformal fluids
[Gubser & Yarom (2010), Staig & Shuryak (2011), Springer & Stephanov (2013)]
- More detailed investigation of linear perturbations and first steps towards non-linear perturbations around Gubser solution
[Hatta, Noronha, Torrieri, Xiao (2014)]
- Linear perturbations around general azimuthally symmetric initial state, realistic equation of state
[Floerchinger & Wiedemann (2013)]
- Characterization of initial conditions by Bessel-Fourier expansion
[Coleman-Smith, Petersen & Wolpert (2012), Floerchinger & Wiedemann (2013)]
- Comparison to full numerical solution shows good convergence properties of perturbative expansion
[Floerchinger, Wiedemann, Beraudo, Del Zanna, Inghirami, Rolando (2013)]
- Related response formalism for expansion in eccentricities
[Teaney & Yan (2012), Yan & Ollitrault (2015)]