

The muon g-2 \longleftrightarrow Δa connection


Massimo Passera
INFN Padova

Max-Planck-Institut für Kernphysik
Heidelberg
16.11.2020

- ➊ Muon g-2: recent theory progress
- ➋ Muon g-2 \iff $\Delta\alpha$ connection
- ➌ The MUonE project

Muon g-2: experimental status

μ



- **BNL 821:** $a_\mu^{\text{EXP}} = (116592089 \pm 54_{\text{stat}} \pm 33_{\text{sys}}) \times 10^{-11}$ [0.5 ppm].
- **Fermilab E989:** new muon g-2 experiment aims at $\pm 16 \times 10^{-11}$ 0.14 ppm. First 3 data taking completed. Analysis of run 1 (~1xBNL) in progress. First result expected very soon with ~ BNL precision.
- **J-PARC:** Muon g-2 proposal. Phase-1 with ~ BNL precision.

Muon g-2: recent theory progress

White Paper of the Muon g-2 Theory Initiative:
arXiv:2006.04822

Muon g-2: the QED contribution

$$a_\mu^{\text{QED}} = (1/2)(\alpha/\pi) \quad \text{Schwinger 1948}$$

$$+ 0.765857426(16) (\alpha/\pi)^2$$

Sommerfield; Petermann; Suura&Wichmann '57; Elend '66; MP '04

$$+ 24.05050988(28) (\alpha/\pi)^3$$

Remiddi, Laporta, Barbieri ... ; Czarnecki, Skrzypek; MP '04;
Friot, Greynat & de Rafael '05, Mohr, Taylor & Newell 2012

$$+ 130.8780(60) (\alpha/\pi)^4$$

Kinoshita & Lindquist '81, ... , Kinoshita & Nio '04, '05;
Aoyama, Hayakawa, Kinoshita & Nio, 2007, Kinoshita et al. 2012 & 2015;
Steinhauser et al. 2013, 2015 & 2016 (all electron & τ loops, analytic);
Laporta, PLB 2017 (mass independent term). **COMPLETED²!**

$$+ 750.86(88) (\alpha/\pi)^5 \text{ COMPLETED!}$$






Kinoshita et al. '90, Yelkhovsky, Milstein, Starshenko, Laporta,...
Aoyama, Hayakawa, Kinoshita, Nio 2012, 2015, 2017 & 2019.
Volkov 1909.08015: A₁⁽¹⁰⁾[no lept loops] at variance, but negligible Δ.

Adding up, we get:


$$a_\mu^{\text{QED}} = 116584718.931(19)(100)(23) \times 10^{-11}$$

from 4-loop & 5-loop coeffs unc. ↪ 6-loop ↪ from α(Cs)

$$a = 1/137.035999046(27) [0.2\text{ppb}] \quad 2018$$



- One-loop term:



$$a_\mu^{\text{EW}}(\text{1-loop}) = \frac{5G_\mu m_\mu^2}{24\sqrt{2}\pi^2} \left[1 + \frac{1}{5} (1 - 4 \sin^2 \theta_W)^2 + O\left(\frac{m_\mu^2}{M_{Z,W,H}^2}\right) \right] \approx 195 \times 10^{-11}$$

1972: Jackiw, Weinberg; Bars, Yoshimura; Altarelli, Cabibbo, Maiani; Bardeen, Gastmans, Lautrup; Fujikawa, Lee, Sanda;
Studenikin et al. '80s


- One-loop plus higher-order terms:

$a_\mu^{\text{EW}} = 153.6 (1.0) \times 10^{-11}$

Hadronic loop uncertainties (and 3-loop nonleading logs).



Muon g-2 TI WP: arXiv:2006.04822

Kukhto et al. '92; Czarnecki, Krause, Marciano '95; Knecht, Peris, Perrottet, de Rafael '02; Czarnecki, Marciano and Vainshtein '02; Degrassi and Giudice '98; Heinemeyer, Stockinger, Weiglein '04; Gribouk and Czarnecki '05; Vainshtein '03; Gnendiger, Stockinger, Stockinger-Kim 2013, Ishikawa, Nakazawa, Yasui, 2019.



The hadronic LO contribution

μ



Keshavarzi, Nomura, Teubner 2018

$$K(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + (1-x)(s/m^2)}$$

$$a_\mu^{\text{HLO}} = \frac{1}{4\pi^3} \int_{4m_\pi^2}^\infty ds K(s) \sigma^{(0)}(s) = \frac{\alpha^2}{3\pi^2} \int_{4m_\pi^2}^\infty \frac{ds}{s} K(s) R(s)$$

$a_\mu^{\text{HLO}} = 6895 (33) \times 10^{-11}$

F. Jegerlehner, arXiv:1711.06089

$= 6939 (40) \times 10^{-11}$

Davier, Hoecker, Malaescu, Zhang, arXiv:1908.00921

$= 6928 (24) \times 10^{-11}$

Keshavarzi, Nomura, Teubner, arXiv:1911.00367


$= 6931 (40) \times 10^{-11}$

Muon g-2 TI WP: arXiv:2006.04822

- Radiative Corrections to $\sigma(s)$ are crucial. S. Actis et al, Eur. Phys. J. C66 (2010) 585
- Great progress in lattice QCD results. Recent BMW result with subpercent precision: $a_\mu^{\text{HLO}} = 7087(53)\times 10^{-11}$. Tension with dispersive evaluations. S. Borsanyi et al. 2002.12347.


The low-energy hadronic cross section

μ



$$R(s) = \sigma(e^+e^- \rightarrow \text{hadrons}) / \frac{4\pi\alpha(s)^2}{3s}$$

Keshavarzi, Nomura Teubner, PRD 2018




Fred Jegerlehner 1905.05078

The hadronic HO VP contribution

μ


- $O(\alpha^3)$ contributions of diagrams containing HVP insertions:



$$a_\mu^{\text{HNLO(vp)}} = -98.3(7) \times 10^{-11}$$

Krause '96; Keshavarzi, Nomura, Teubner 2019; Muon g-2 TI WP.

- $O(\alpha^4)$ contributions of diagrams containing HVP insertions:




$$a_\mu^{\text{HNNLO(vp)}} = 12.4(1) \times 10^{-11}$$

Kurz, Liu, Marquard, Steinhauser 2014

- HNLO light-by-light

⌚ This term had a troubled life! Nowadays:




$a_\mu^{\text{HNLO}}(bl) = + 80 (40) \times 10^{-11}$	Knecht & Nyffeler '02
$= +136 (25) \times 10^{-11}$	Melnikov & Vainshtein '03
$= +105 (26) \times 10^{-11}$	Prades, de Rafael, Vainshtein '09
$= + 100 (29) \times 10^{-11}$	Jegerlehner, arXiv:1705.00263
$= + 92 (19) \times 10^{-11}$	Muon g-2 TI WP, 2006.04822

- ⌚ Significant improvements due to data-driven dispersive approach.
⌚ Great progress on the lattice. Recent RBC result: $79(35)\times 10^{-11}$ arXiv:1911.08123

- HNNLO light-by-light

$$a_\mu^{\text{HNNLO}}(|bl|) = 2 (1) \times 10^{-11}$$

Colangelo, Hoferichter, Nyffeler, MP, Stoffer 2014; Muon g-2 TI WP, 2006.04822



Comparing the SM prediction with the measured muon g-2 value:

$$a_\mu^{\text{EXP}} = 116592089 (63) \times 10^{-11}$$

BNL E821

$$a_\mu^{\text{SM}} = 116591810 (43) \times 10^{-11}$$

Muon g-2 TI

$$\Delta a_\mu = a_\mu^{\text{EXP}} - a_\mu^{\text{SM}} = 279 (76) \times 10^{-11}$$

3.7 σ

Muon g-2 \iff $\Delta\alpha$ connection

Marciano, MP, Sirlin 2008 & 2010

Keshavarzi, Marciano, MP, Sirlin 2020

- Can Δa_μ be due to missing contributions in the hadronic $\sigma(s)$?
- An upward shift of $\sigma(s)$ also induces an increase of $\Delta a_{\text{had}}^{(5)}(M_Z)$.
- Consider:

$$\begin{aligned} a_{\mu^{\text{HLO}}} &\rightarrow a = \int_{4m_\pi^2}^{s_u} ds f(s) \sigma(s), \quad f(s) = \frac{K(s)}{4\pi^3}, \quad s_u < M_Z^2, \\ \Delta a_{\text{had}}^{(5)} &\rightarrow b = \int_{4m_\pi^2}^{s_u} ds g(s) \sigma(s), \quad g(s) = \frac{M_Z^2}{(M_Z^2 - s)(4\alpha\pi^2)}, \end{aligned}$$

and the increase


$$\Delta\sigma(s) = \epsilon\sigma(s)$$

$\epsilon > 0$, in the range:

$$\sqrt{s} \in [\sqrt{s_0} - \delta/2, \sqrt{s_0} + \delta/2]$$



How much does the M_H upper bound from the EW fit change when we shift up $\sigma(s)$ by $\Delta\sigma(s)$ [and thus $\Delta\alpha_{\text{had}}^{(5)}(M_Z)$] to accommodate $\Delta\alpha_\mu$?




Marciano, MP, Sirlin, 2008 & 2010


Major update: Higgs discovered, improved EW observables (M_W , $\sin^2\theta$, M_{top} , ...), updates to $\sigma(s)$, theory improvements, global fit, ...

Parameter	Input value	Reference	Fit result	Result w/o input value
M_W (GeV)	80.379(12)	[5]	80.359(3)	80.357(4)(5)
M_H (GeV)	125.10(14)	[5]	125.10(14)	94^{+20+6}_{-18-6}
$\Delta\alpha_{had}^{(5)}(M_Z^2) \times 10^4$	276.1(1.1)	[23]	275.8(1.1)	272.2(3.9)(1.2)
m_t (GeV)	172.9(4)	[5]	173.0(4)	...
$\alpha_s(M_Z^2)$	0.1179(10)	[5]	0.1180(7)	...
M_Z (GeV)	91.1876(21)	[5]	91.1883(20)	...
Γ_Z (GeV)	2.4952(23)	[5]	2.4940(4)	...
Γ_W (GeV)	2.085(42)	[5]	2.0903(4)	...
σ_{had}^0 (nb)	41.541(37)	[108]	41.490(4)	...
R_l^0	20.767(25)	[108]	20.732(4)	...
R_c^0	0.1721(30)	[108]	0.17222(8)	...
R_b^0	0.21629(66)	[108]	0.21581(8)	...
\bar{m}_c (GeV)	1.27(2)	[5]	1.27(2)	...
\bar{m}_b (GeV)	$4.18^{+0.03}_{-0.02}$	[5]	$4.18^{+0.03}_{-0.02}$...
$A_{FB}^{0,l}$	0.0171(10)	[108]	0.01622(7)	...
$A_{FB}^{0,c}$	0.0707(35)	[108]	0.0737(2)	...
$A_{FB}^{0,b}$	0.0992(16)	[108]	0.1031(2)	...
A_ℓ	0.1499(18)	[75,108]	0.1471(3)	...
A_c	0.670(27)	[108]	0.6679(2)	...
A_b	0.923(20)	[108]	0.93462(7)	...
$\sin^2\theta_{eff}^{lep}(Q_{FB})$	0.2324(12)	[108]	0.23152(4)	0.23152(4)(4)
$\sin^2\theta_{eff}^{lep}(\text{Had Coll})$	0.23140(23)	[100]	0.23152(4)	0.23152(4)(4)

Keshavarzi, Marciano, MP, Sirlin, PRD 2020 (using Gfitter)



Shifts $\Delta\sigma(s)$ to fix Δa_μ are possible,
but conflict with the EW fit if they occur above ~ 1 GeV




Uniform scaling of $\sigma(s)$ below $\sim 0.7 \text{ GeV}$? +9% required!

Keshavarzi, Marciano, MP, Sirlin, PRD 2020

How large are the required shifts $\Delta\sigma(s)$?

$\Delta\alpha$



Shifts below 1 GeV conflict with the quoted exp. precision of $\sigma(s)$

Keshavarzi, Marciano, MP, Sirlin, PRD 2020

What happens to the electron g-2?

- The 2008 measurement of the electron g-2 is:

$$a_e^{\text{EXP}} = 11596521807.3 (2.8) \times 10^{-13} \quad \text{Hanneke et al, PRL100 (2008) 120801}$$

vs. old (factor of 15 improvement, 1.8σ difference):

$$a_e^{\text{EXP}} = 11596521883 (42) \times 10^{-13} \quad \text{Van Dyck et al, PRL59 (1987) 26}$$

- Equate $a_e^{\text{SM}}(\alpha) = a_e^{\text{EXP}}$ → “g_e-2” determination of alpha:

$$\alpha^{-1} = 137.035\ 999\ 151 (33) \quad [0.24 \text{ ppb}]$$

- The best determination of α is obtained via atomic interferometry:

$$\alpha^{-1} = 137.036\ 999\ 046 (27) \quad [0.20 \text{ ppb}] \quad \text{Science 360 (2018) 191 (Cs)}$$

(was $\alpha^{-1} = 137.035\ 998\ 995 (85)$ [0.62 ppb] PRL106 (2011) & CODATA 2016)

2.5 sigma discrepancy

Electron g-2: SM vs experiment

- Using $\alpha = 1/137.036\ 999\ 046\ (27)$ [Cs 2018], the SM prediction for the electron g-2 is:

$$a_e^{\text{SM}} = 115\ 965\ 218\ 16.2\ (0.1)\ (0.1)\ (2.3) \times 10^{-13}$$

δC_5^{qed} δa_e^{had} from $\delta \alpha$

- The (EXP – SM) difference is:

$$\Delta a_e = a_e^{\text{EXP}} - a_e^{\text{SM}} = -8.9\ (3.6) \times 10^{-13} [2.5\sigma]$$

NB: negative!

Note the negative sign. [QED 5-loop $a_e^{\text{QED5}} = 4.6 \times 10^{-13}$]

- NP sensitivity limited only by the experimental errors in α and a_e . May soon play a pivotal role in probing NP in the leptonic sector.

Giudice, Paradisi, MP 2012

- The present sensitivity is $\delta\Delta a_e = 3.6 \times 10^{-13}$, ie (10^{-13} units):

$$(0.1)_{\text{QED5}}, \quad (0.1)_{\text{HAD}}, \quad (2.3)_{\delta\alpha}, \quad (2.8)_{\delta a_e^{\text{EXP}}}$$

$\overbrace{\qquad\qquad\qquad}^{(0.1)_{\text{TH}}}$

- The $(g-2)_e$ exp. error may soon drop below 10^{-13} and work is in progress to further reduce the error induced by $\delta a \rightarrow$

sensitivity below 10^{-13} may be reached with ongoing exp work

- In a broad class of BSM theories, contributions to a_l scale as

$$\frac{\Delta a_{\ell_i}}{\Delta a_{\ell_j}} = \left(\frac{m_{\ell_i}}{m_{\ell_j}} \right)^2$$


This **Naive Scaling** leads to:

$$\Delta a_e = \left(\frac{\Delta a_\mu}{3 \times 10^{-9}} \right) 0.7 \times 10^{-13}; \quad \Delta a_\tau = \left(\frac{\Delta a_\mu}{3 \times 10^{-9}} \right) 0.8 \times 10^{-6}$$

Giudice, Paradisi & MP, JHEP 2012

Shift of the electron g-2

e




Shifts $\Delta\sigma(s)$ to fix Δa_μ slightly increase the $|\Delta a_e| \sim 10^{-12}$ tension

Keshavarzi, Marciano, MP, Sirlin, PRD 2020

Shift of the e/μ g-2 scaled HLO ratio

e/μ



**Good agreement between lattice [Giusti & Simula 2020] and KNT19.
Possible future bounds on very low energy shifts $\Delta\sigma(s)$?**


Keshavarzi, Marciano, MP, Sirlin, PRD 2020

- Crivellin, Hoferichter, Manzari and Montull, “Hadronic vacuum polarization: $(g-2)_\mu$ versus global electroweak fits,” PRL125 (2020) 091801 [arXiv:2003.04886].
- Eduardo de Rafael, “On Constraints Between $\Delta\alpha_{\text{had}}(M^2)$ and $(g_\mu-2)_{\text{HVP}}$,” arXiv:2006.13880.
- Malaescu and Schott, “Impact of correlations between a_μ and a_{QED} on the EW fit”, arXiv:2008.08107.
- Colangelo, Hoferichter and Stoffer, “Constraints on the two-pion contribution to hadronic vacuum polarization,” arXiv:2010.07943.

The MUonE project




- The leading hadronic contribution a_μ^{HLO} computed via the timelike formula:



$$a_\mu^{\text{HLO}} = \frac{1}{4\pi^3} \int_{4m_\pi^2}^\infty ds K(s) \sigma_{\text{had}}^0(s)$$

$$K(s) = \int_0^1 dx \frac{x^2 (1-x)}{x^2 + (1-x) (s/m_\mu^2)}$$

- Alternatively, simply exchanging the x and s integrations:




$$a_\mu^{\text{HLO}} = \frac{\alpha}{\pi} \int_0^1 dx (1-x) \Delta\alpha_{\text{had}}[t(x)]$$

$$t(x) = \frac{x^2 m_\mu^2}{x-1} < 0$$

Lautrup, Peterman, de Rafael, 1972

$\Delta\alpha_{\text{had}}(t)$ is the hadronic contribution to the running of α in the spacelike region: a_μ^{HLO} can be extracted from scattering data!



- $\Delta\alpha_{had}(t)$ can be measured via the **elastic scattering** $\mu e \rightarrow \mu e$.
- We propose to scatter a 150 GeV muon beam, available at CERN's North Area, on a fixed electron target (Beryllium). Modular apparatus: each station has one layer of Beryllium (target) followed by several thin Silicon strip detectors.



Abbiendi, Carloni Calame, Marconi, Matteuzzi, Montagna,
Nicrosini, MP, Piccinini, Tenchini, Trentadue, Venanzoni


EPJC 2017 - arXiv:1609.08987

- For a 150 GeV muon beam, MUonE's scan region extends up to $x=0.932$, ie beyond the peak! (the peak is at $x=0.914$)
- The high-energy region inaccessible to MUonE contributes only 13% of a_μ^{HLO} integral. It can be determined with timelike data and/or lattice QCD results. Recently obtained via lattice QCD. Giusti&Simula and Marinkovic'&Cardoso 2019



- **Statistics:** With CERN's 150 GeV muon beam M2 ($1.3 \times 10^7 \mu/\text{s}$), incident on 40 15mm Be targets (total thickness 60cm), 2-3 years of data taking ($2 \times 10^7 \text{ s/yr}$) $\rightarrow \mathcal{L}_{\text{int}} \sim 1.5 \times 10^7 \text{ nb}^{-1}$.
- With this \mathcal{L}_{int} we estimate that measuring the shape of $d\sigma/dt$ we can reach a statistical sensitivity of $\sim 0.3\%$ on a_{μ}^{HLO} , ie $\sim 20 \times 10^{-11}$.
- **Systematic** effects must be known at $\lesssim 10\text{ppm}$!
- Test beams performed at CERN in 2017 & 2018 [arXiv:1905.11677](#)
- Lol submitted to CERN SPSC in 2019: Test run approved for 2021.
- Full-statistics run hopefully in 2022–24.

- To extract $\Delta\alpha_{\text{had}}(t)$ from MUonE's measurement, the ratio of the SM cross sections in the signal and normalisation regions must be known at $\lesssim 10\text{ppm}$!



- Fully differential fixed-order MC @ NLO ready Pavia and PSI 2018-19
- NNLO QED: MI for 2-loop box diagrams computed Padova 2017-19
- Two MC built including partial subsets of the NNLO QED corrections due to electron and muon radiation Pavia and PSI 2020
- NNLO hadronic effects computed Padova and Siegen 2019
- Extraction of the leading electron mass effects from the massless muon-electron scattering amplitudes PSI 2019-2020
- ...

Theory for muon-electron scattering @ 10 ppm:
A report of the MUonE theory initiative. arXiv:2004.13663

- MUonE will be very precise → could NP affect its measurements?
Using existing experimental bounds we showed that this is **unlikely**:
- Consider “light” or “heavy” mediators [ie, mass lower or higher than MUonE’s energy scale of $O(1 \text{ GeV})$]:
 - Heavy NP — EFT formalism:
 - S & T effects suppressed by electron mass and, for T, also by $(g-2)_e$.
 - P doesn’t interfere with QED
 - V & A effects excluded by $e^+e^- \rightarrow \mu^+\mu^-$ data (mainly).
 - LFV effects excluded by muonium-antimuonium oscillation limits.
 - Light NP — spin 0 and 1 mediators:
 - ALPs, Dark Photons and light Z’ bosons effects excluded by direct searches and dipole moments. LVF excluded by muonium oscillation.
- NB: MUonE will not be sensitive to NP signals which are constant (in t) relative to the LO QED one!

Dev, Rodejohann, Xu, Zhang, JHEP 2020
 Masiero, Paradisi, MP, PRD 2020

MUonE — Theory workshops



Padova

Europe/Rome timezone

Muon-electron scattering: Theory kickoff workshop

4-5 September 2017

Overview


Venue

Timetable

Logistic

Map

 Support



**MUonE theory workshops: Padova 2017, Mainz 2018, Zurich 2019
Next MUonE theory workshop: MITP Mainz 1-5.03 2021 (hopefully)**



Conclusions

- FNAL's first a_μ result expected very soon with \sim BNL precision.
- Is present Δa_μ due to missed contributions in the hadronic $\sigma(s)$?
Shifts $\Delta\sigma(s)$ to fix Δa_μ conflict with the global EW fit above ~ 1 GeV
Shifts below ~ 1 GeV conflict with the quoted exp. error of $\sigma(s)$.
- Shifts $\Delta\sigma(s)$ to fix Δa_μ slightly increase the a_e tension ($R_{e/\mu}$ ok).
- MUonE at CERN will provide a new independent spacelike determination of a_μ^{HLO} alternative to the DR and lattice ones.


Backup

Muon g-2: connection with M_w and $\sin^2\theta$


$\Delta\alpha$



Keshavarzi, Marciano, MP, Sirlin, PRD 2020



Keshavarzi, Marciano, MP, Sirlin, PRD 2020



Keshavarzi, Marciano, MP, Sirlin, PRD 2020