

# Exercises to “Standard Model of Particle Physics II”

Winter 2015/16

Prof. Dr. Manfred Lindner, Dr. Werner Rodejohann

Sheet 10

13.01.16

## **Exercise 21:** Left-right symmetric electroweak model I [15 Points]

The left-right symmetric model can be introduced by assuming right-handed fermion doublets in analogy to the left-handed ones. The quark and lepton spectra consist of

$$Q_{L,R}^i = \begin{pmatrix} U_{L,R}^i \\ D_{L,R}^i \end{pmatrix}; \quad L_{L,R}^i = \begin{pmatrix} \nu_{L,R}^i \\ e_{L,R}^i \end{pmatrix},$$

with the following  $SU(2)_L$ ,  $SU(2)_R$ ,  $U(1)_{B-L}$  transformation properties:

$$\begin{aligned} Q_L &: (2_L, 1_R, 1/3); & L_L &: (2_L, 1_R, -1); \\ Q_R &: (1_L, 2_R, 1/3); & L_R &: (1_L, 2_R, -1). \end{aligned}$$

The Higgs sector contains a bi-doublet  $\phi$  and two triplets  $\Delta_L$  and  $\Delta_R$  with transformation properties

$$\phi : (2_L, 2_R, 0); \quad \Delta_L : (3_L, 1_R, 2); \quad \Delta_R : (1_L, 3_R, 2).$$

Note that the bi-doublet and the triplets can be expressed by the following  $2 \times 2$  matrices:

$$\phi = \begin{pmatrix} \phi_1^0 & \phi_2^+ \\ \phi_1^- & \phi_2^0 \end{pmatrix}; \quad \Delta_{L,R} = \begin{pmatrix} \Delta^+/\sqrt{2} & \Delta^{++} \\ \Delta^0 & -\Delta^+/\sqrt{2} \end{pmatrix}.$$

- Why does this model not work with only the bi-doublet?
- Construct the Lagrange density for the fermion-Higgs interactions  $\mathcal{L}_{\text{Yukawa}}$ .
- Use the assumption that, after spontaneous symmetry breaking the vacuum is electrically neutral, write down the fermion mass terms in the broken phase.
- We leave the quark and lepton sector unchanged, but modify the symmetry-breaking part of the model. In the Higgs sector we still have the bi-doublet  $\phi$ , but instead of the triplets we introduce two scalar doublets  $A_{L,R}$  and a fermionic singlet  $\chi$  with the following transformation properties under  $SU(2)_L$ ,  $SU(2)_R$  and  $U(1)_{B-L}$ :

$$A_L : (2_L, 1_R, 1); \quad A_R : (1_L, 2_R, 1); \quad \chi : (1_L, 1_R, 0).$$

Left-right symmetry implies the invariance of the Lagrange density under the following transformations (where  $\Psi$  denotes any fermion field):

$$\Psi_L \leftrightarrow \Psi_R \quad A_L \leftrightarrow A_R \quad \phi \leftrightarrow \phi^\dagger.$$

Construct the Lagrange density  $\mathcal{L}_{\text{Yukawa}}$  for the fermion masses in this model (you have to build singlets under the whole gauge group).

**Exercise 22:**  $Z'$  physics [15 Points]

The general effective Lagrange density after breaking the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)'$  symmetry to  $SU(3)_C \times U(1)_{EM}$  can be written as

$$\mathcal{L} = \mathcal{L}_{SM} + \mathcal{L}_{Z'} + \mathcal{L}_{mix},$$

where the relevant part of the Standard Model Lagrangian is

$$\mathcal{L}_{SM} = -\frac{1}{4}\hat{B}_{\mu\nu}\hat{B}^{\mu\nu} - \frac{1}{4}\hat{W}_{\mu\nu}^a\hat{W}^{\mu\nu,a} + \frac{1}{2}\hat{M}_Z^2\hat{Z}_\mu\hat{Z}^\mu - \frac{\hat{e}}{\hat{c}_W}j_B^\mu\hat{B}_\mu - \frac{e}{c_W}j_W^{\mu,a}\hat{W}_\mu^a$$

and the hats merely denote that the fields are not mass eigenstates. The  $Z'$  part reads

$$\mathcal{L}_{Z'} = -\frac{1}{4}\hat{Z}'_{\mu\nu}\hat{Z}'^{\mu\nu} + \frac{1}{2}\hat{M}_{Z'}^2\hat{Z}'_\mu\hat{Z}'^\mu - g'j'_\mu\hat{Z}'^\mu$$

and the kinetic- and mass-mixing terms can be parameterized as

$$\mathcal{L}_{mix} = -\frac{\sin\chi}{2}\hat{Z}'_{\mu\nu}\hat{B}^{\mu\nu} + \delta\hat{M}^2 Z'_\mu Z'_\mu \hat{Z}^\mu$$

- Determine the mass eigenstates  $Z_1^\mu$  and  $Z_2^\mu$  and determine the couplings of  $Z_{1,2}$  to the currents  $j_B$ ,  $j_W$  and  $j'$ . Set the kinetic mixing angle  $\chi$  to zero for simplicity.
- Since the mass of the physical  $Z$  boson changes compared to the SM, the  $\rho$  parameter is no longer equal to one (at tree-level). Use the current value  $\rho = 1.0008_{-0.0007}^{+0.0017}$  to constrain the  $Z$ - $Z'$  mixing.
- A well-motivated extension of the SM is a gauged  $B-L$  symmetry (baryon minus lepton-number). Write down explicitly the corresponding current for the SM fermions:

$$j'_\mu = \sum_\psi \bar{\psi} \gamma_\mu (B-L) \psi.$$

- To cancel quantum anomalies in the  $B-L$  model, one also has to introduce three right-handed neutrinos  $N_i$  which are then part of the current:  $\Delta j'_\mu = -\sum_i \bar{N}_i \gamma_\mu P_R N_i$ . Due to  $Z$ - $Z'$  mixing, the SM-like  $Z_1$  will also couple to these new neutrinos. Can you give a constraint on the  $Z$ - $Z'$  mixing from the well-measured  $Z$ -width (for  $M(N_i) \ll M_Z/2$ )? See the PDG for numbers.

**Tutor:**

Miguel Campos, email: miguel.campos@mpi-hd.mpg.de

Lectures homepage: [https://www.mpi-hd.mpg.de/manitop/StandardModel2/index\\_WS15.html](https://www.mpi-hd.mpg.de/manitop/StandardModel2/index_WS15.html)

**Discussion and hand-in of sheet:**

Tuesday 19.01.16 4.15 pm, INF 501/CIP R. 103