

# Exercises to “Standard Model of Particle Physics II”

Winter 2022/23

Prof. Dr. Manfred Lindner and Dr. Werner Rodejohann  
Sheet 07 - November 30, 2022

---

**Tutor:** Sophie Klett   **e-mail:** [sophie.klett@mpi-hd.mpg.de](mailto:sophie.klett@mpi-hd.mpg.de)

**Lecture webpage:** <https://www.mpi-hd.mpg.de/manitop/StandardModel2/index.html>

**Hand-in of solutions:**

December 7, 2022 - 11:15, Phil. 12, R105

**Discussion of solutions:**

December 7, 2022 - 11:15, Phil. 12, R105

---

## Problem 1: *Mass matrices and mixing angles* [15 Points]

A general (Dirac) mass term for fermions is given by

$$\mathcal{L}_M = \bar{\psi}_{i,L} M_{ij} \psi_{j,R} + \text{h.c.}$$

where  $M$  is hermitian and given by a  $n \times n$  Yukawa coupling matrix  $Y$  times the Higgs vev.

- Show that for an arbitrary  $n \times n$  matrix  $M$  one can choose a bi-unitary transformation  $UMV^\dagger$  to diagonalize  $M$ , such that no diagonal element  $UMV^\dagger = D := \text{diag}(m_1, m_2, \dots, m_n)$  is negative. The matrices  $U$  and  $V$  are unitary.
- Show that for a real mass matrix  $M$  one can choose orthogonal diagonalization matrices.
- As an example for calculable mixing angles consider a simple  $2 \times 2$  mass matrix of the form

$$M = \begin{bmatrix} 0 & a \\ a^* & b \end{bmatrix}.$$

The unitary matrix that diagonalizes  $M$  can be described by a single parameter: a *mixing angle*  $\theta$ . Show that the following relation between mixing angle and masses holds:

$$\tan \theta = \sqrt{\frac{m_1}{m_2}}.$$

Compare this with the Cabibbo angle and the down and strange quark masses.

- A completely different situation holds for the symmetric mass matrix

$$M = \begin{bmatrix} a & b & b \\ b & \frac{1}{2}(a+b+d) & \frac{1}{2}(a+b-d) \\ b & \frac{1}{2}(a+b-d) & \frac{1}{2}(a+b+d) \end{bmatrix}.$$

Give the mixing matrix for this mass matrix (*Hint:* try first a 23-rotation).

## Problem 2: *Mixing of leptons* [5 Points]

Show that a mixing matrix for charged leptons would have no physical effect if neutrinos were massless particles. In other words, charged lepton mixing in the Standard Model with massless neutrinos is redundant.